



Assessment of environmental quality in soil under wheat and vines in Bouznika-Benslimane region of Morocco

F. Zaakour*, Z. Chems, F. Matech, K. Moustarhfer, N. Saber*

Université Hassan II -Mohammedia-Casablanca, Faculté des Sciences Ben M'Sik, Département de Géologie, BP 7955
Casablanca, Maroc

Received 7 Dec 2013, Revised 16 July 2014, Accepted 17 July 2014

*Corresponding Authors. E-mail: Zaakour-fatna@hotmail.com; Tel:(+212606438139) & najib_saber@yahoo.fr; Tel: (+212667900025)

Abstract

This work aims at assessing the environmental quality of agricultural soils under vine and wheat cultivation in the area of Bouznika-Benslimane (Morocco). 50 soil samples were collected and studied. The results obtained been studied and recorded on the agronomic parameters, show that the soil's texture is a little clay with a slightly acid pH neutral. Moreover, the electrical conductivity values are clearly higher in soils under wheat than in those under vine. The variability of CaCO₃ contents within the different studied soils is generally similar in all stations. Moreover, the organic substance rate seems to be more important in the soils under the wheat crops (2%) than in those under the vine crops (1.7%). The examined soil samples under vine and wheat have shown that these soils contain an important concentration of metal trace elements, essentially the Cu (0.81 ppm), the Cd (0.85 ppm), the Ni (81.86 ppm), the Pb (67.14 ppm) and the Zn (791.63 ppm). These metallic elements were also detected in significant concentrations within the soils under vine with a pollution index of high (PI).

Keywords: Bouznika-Benslimane, Agricultural soils, Vines, Wheat, Agronomic parameters, Heavy metals.

1.Introduction

When considering different kinds of contaminants, heavy metals are especially dangerous because of their persistence and toxicity [1]. Heavy metal accumulation in soils can result in a loss of soil functions leading to concerns about environmental quality protection, maintenance of human health and productivity. Soil pollution can have implications in phytotoxicity at high concentrations and result in the transfer of heavy metals to the human diet from crop uptake or soil ingestion by grazing livestock [2-5].

There are two main sources of heavy metals in the soil [6]: (i) natural background, which represents the heavy metal concentration derived from parent rocks; (ii) including agrochemicals, organic amendments, animal manure, mineral fertilizer, sewage sludge and industrial wastes. In the last several decades, the natural input of several heavy metals to soils due to pedogenesis has been exceeded by the human input, even on global and regional scales [7].

The importance of pH in metal solubility is well-known as it influences heavy metal adsorption, retention and movement [8-9]. Related to this, Matos and al have demonstrated that pH and Ca concentration affect Cd, Cu, Pb and Zn mobility in soils by competition of adsorbing sites [10].

Organic matter is another important soil component that influences metal availability. It has a nutritional function by serving as a source of N, P and S, and has a high binding capacity for cations and organic contaminants [11].The heavy metals are accumulated in surface organic layers in agricultural and urban soils [12-14].

The aim of this study is to evaluate the state of the heavy metal content in agricultural soils in relation wheat and vines in the region of Bouznika-Benslimane (Morocco). This evaluation takes into account the physical and chemical aspects of the soil.

2. Materials and methods

2.1. Field of study

The area of this study belongs to the provinces of the Bouznika and Benslimane cities all of which are the Atlantic coast of Morocco. This area is between the cities of Rabat and Casablanca (Figure1). This region is characterized by a semi-arid

Mediterranean climate. The annual rainfall average of the last 20 years in the region is about 400 mm. Oceanic influence creates variations at the level of the forest microclimate. These changes result in an increase of the coast rainfall in Benslimane. In other words, the rainfall tends to decrease by the North-South gradient. The minimum and maximum annual temperature average is respectively 10.3 and 23.7 °C [15].




Figure 1: Location of study area

The metal contamination degree of soil is often estimated through the total contents of heavy metals in the surface horizon [16]. Throughout this work, five stations within the region of Bouznika Benslimane are our main field of study. We have looked at 10 samples in each station in the horizon [0-20 cm]: (5 under the cultivation of wheat and 5 under vine).

The stations are selected and classified according to the treatment type of the vine cultivation. Those stations are namely Ouled Taleb and Cherrate, stations which are characterized by a traditional treatment. Then, there are the stations of Skhairate and Mansfield which are modern stations. The last one is Bouznika which is considered as a check station.

2.2. The analysis method

The analysis of the texture in the various stations determined by the method of Robinson pipette [17], the analysis of the organic substance has been conducted in the method of Walkley and Black [18], the electrical conductivity in the method of Rhoades [19], the pH in the method of Mc Lean [20] while the CaCO₃ dosing was made by the method of Bernard [21].

The determination of heavy metals is carried out by ICP-AES method at the National Centre for Scientific and Technical Research (Rabat). The analytical laboratory follows the standards (1000 ppm accurate JobinYvon) certified by the ISO 9001 quality assurance system.

2.3. Statistical analyzes

Statistical analyzes were performed with the help of the XLSTAT 2013 statistical software. The analysis of variance was used to find the significant effect of environmental parameters (Cd, Cu, Fe, Ni, Zn, pb and Al) and of the other plots.

3. Results and discussion

3.1. Agronomic parameters

The analysis of texture in the various stations revealed that the texture is clay in the Ouled Taleb, silty clay loam in Bouznika and Mansouria, silt loam in Skhairate and clay loam in Cherrate station (Figure 2).

The pH is considered the main chemical parameter controlling the bioavailability of heavy metals in the soil [22]. The results of this study show that the pH is neutral to slightly acid in all stations. The pH values are higher in Skhairate and Ouled Taleb (Figure3.a).

The measurements of the electrical conductivity are lower in the soils stations of Bouznika, Mansouria and Ouled Taleb than in the soils of Skhairate and Cherrate. Measurements are higher in soils under wheat cultivation in comparison with that of soil under vine (excepted Mansouria station) (Figure3.b).

The CaCO₃ values are similar under wheat and under vine in Bouznika, Skhairate and Ouled Taleb stations, however they can vary in the stations of Cherrate and Mansouria (Figure3.c).

Moreover, the average rate of organic substance in the wheat plots is higher (2%) than under vine (1.7%) with the exception of the plot of land under wheat in Skhairate where there is a supply of organic fertilizer (Figure3.d).




Figure 2: Texture of soils in areas of Bouznika Benslimane




Figure 3: Variations in pH, EC, CaCO₃ and soil organic matter under vine and under wheat stations studied

3.2. Environmental parameters

The concentration of metal trace elements (MTE) that is measured in the soil is the outcome of the heritage created by the alteration of the rock concentrations (local geochemical background), more or less altered by soil formation process (soil geochemical background). This background is enhanced by the contributions of human activities [23].

The aluminium contents of soils under wheat are high in comparison with soils under vine in Cherrate station with an important difference between soils under vine and soils under wheat. (Figure4.a). The concentrations of Fe, which are the lowest, are recorded in the stations of Mansouria (6.91 ppm) and of Bouznika (9.49 ppm) while the maximum concentrations are obtained in the soils of the region of Ouled Taleb (65.31 ppm). The difference in the iron concentrations between plots under vine and plots under wheat is highly significant in Cherrate and Skhairate stations (Figure.4.b).

The maximum cadmium was obtained in the under vine soil within the area of Cherrate with a grade of 0.81 ppm. The lowest cadmium content was observed in the Skhairate station with an average value of 0.52 ppm. The difference of concentrations is highly significant between plots under vine and plots under wheat at the Ouled Taleb, Cherrate and Skhairate stations (Figure.4.c).

Fungicide treatments of the vines have long been based primarily on the salt spray of copper (Cu) on the vine leaves. Their widespread use since the late 19th century has led to the pollution of Cu that resulted in risks such as the Cu phytotoxicity which threatens the plants grown in these soils [24]. In our study, the soil under vine shows high or very high levels of Cu in comparison with the one under wheat (except Mansouria and Ouled Taleb stations). The maximum concentrations of Cu were recorded in soils of Ouled Taleb (66.70 ppm) and Bouznika for Wheat (0.92 ppm), the difference in copper concentrations between stations is highly significant (Figure4.d).

The highest concentrations of Ni are recorded in the Ouled Taleb under wheat soils. These ones were cultivated with vines before wheat. The difference of Nickel concentrations between plots under vine and wheat is highly significant in all stations (Figure4.e).

The levels of lead found in all stations under study do not generally exceed the normal value of uncontaminated soils (100 ppm) set by the OMS. Thus, the maximum lead is recorded in the under vineyards soils which are treated in the traditional way (from 40.84 to 67.14 ppm), while areas of Mansouria and Bouznika show low levels of lead with an average of (38.36 to 40.84 ppm). The difference in lead concentrations between plots of vine and wheat is highly significant in all stations (Figure4.f).

The Zinc substance which is an essential element is to some extent low. However, it causes an acute toxicity with a high dose [25]. The average concentrations in soil under vines are greater than those obtained in the soil under wheat in all the stations with the exception of Ouled Taleb station. The difference in zinc concentrations between the plots of vine and wheat is highly significant in the stations of Cherrate and Mansouria (Figure4.g).




Figure 4: Soil Al, soil Fe, soil Cd, soil Cu, soil Ni, soil Pb and soil Zn

The average concentrations of the Cu, Cd, Ni, and Pb found in the soils of the studied regions are below the OMS standard which is namely 2 ppm Cd, 100 ppm Cu, 50 ppm Ni, and 100 ppm for pb [26].

The average concentrations in soil under vine Zn are above the standard set by the OMS (300 ppm) in the stations of Bouznika, Shkairate and OuledTaleb while they exceed it in the of stations of Cherrate (274 to 378.35 ppm) and Mansouria (261.98 to 791.63 ppm).

In addition, very low levels of iron (6.91 to 65.3 ppm) are recorded in the soils being studied at the level of all stations in comparison with the average maximum levels recommended by the OMS (10 000- 50 000 ppm). These values have allowed us to detect iron deficiency in our soils.

The correlation test between the different parameters being studied allowed us to highlight. Positive correlations were found between pH, EC and organic matter while moderate correlations were observed between Cadmium and Nickel, Lead and Zinc (Table 1).

Table1: The best correlation coefficients obtained in this study.

	pH	CaCo ₃	MO	CE	Cd	Cu	Fe	Ni	Pb	Zn	Al
pH	1,000										
CaCo ₃	-0,292	1,000									
MO	0,715	-0,526	1,000								
CE	0,773	-0,076	0,637	1,000							
Cd	-0,423	0,149	-0,186	-0,246	1,000						
Cu	0,103	-0,111	0,300	-0,092	0,263	1,000					
Fe	0,172	-0,146	0,380	0,000	0,273	0,989	1,000				
Ni	-0,135	-0,012	-0,004	-0,253	0,429	0,322	0,331	1,000			
Pb	-0,227	0,232	-0,105	-0,128	0,425	0,057	0,028	0,014	1,000		
Zn	-0,538	0,507	-0,716	-0,239	0,480	-0,411	-0,417	0,063	0,246	1,000	
Al	0,325	0,039	0,082	0,304	0,277	-0,278	-0,233	0,406	0,195	0,257	1,000

Iron concentrations in the different studied soils are perfectly correlated with copper concentrations (Figure 5). Negative and remarkable correlations are recorded between the Zinc, pH, organic matter, copper and iron. In addition, positive correlations were found between the CaCO₃ and Zinc, Cadmium and Lead and between Nickel and Cadmium, Copper and Iron (Table 1).




Figure 5: Diagram Correlation iron / copper in soils studied

The heavy metals contamination in the environmental surface is related to all the contaminants rather than to one metal. Several authors have introduced the pollution index of soil to identify the multi-element contamination which leads to an increase in the metal pollution [27-29].

The pollution index is defined as the rappers means of the metal concentrations in soil samples against the limit values [30].

The calculated values of pollution index shows that soil under vines are more polluted soil under wheat. This pollution seems to reduce in the control station of Bouznika.

The decrease in pollution in the wheat plots of land is comparable to other wheat plots with the exception of the OuledTaleb station.

In fact, in this station, the soil under wheat show high concentrations of Cd (0.724 ppm), Cu (54.696 ppm), Fe (59.868 ppm), Ni (44.762 ppm), Pb and 54.338 ppm (Figure6).

The spatial variation of the pollution index in the region of Bouznika-Benslimane indicates maximum values observed in soil under vine station Mansouria (0.508) while the lowest values are noticed in the under wheat soils within the Skhairate area (0.298).




Figure 6: Indices of pollution soils under vine and wheat in each station

Conclusion

The agronomical characterization of the study of soil has allowed us to demonstrate that the electrical conductivity, pH and organic matter are significantly higher in soils under wheat compared with soils under vine.

The contamination evaluation in the soils under vine and wheat within the region of Bouznika-Benslimane and by the metal trace elements (Al, Cd, Cu, Fe, Ni, and Pb) shows that these soils have concentrations below the established standards OMS, with an abnormality detected for Zn especially for soils under vine. Indeed, some values of this element exceed the critical threshold of 300 ppm set by OMS. The Iron deficiency was recorded at the five study sites.

From a statistical point of view we could highlight:

- positive correlations between organic matter, pH and EC;
- a perfect correlation between Cu and Fe;
- strong correlations between the Zn, the pH, the CaCO₃, the organic matter, the Cd, the Cu, and the Fe.

The pollution index, calculated in the layer [0-20 cm], shows higher values under vine in the OuledTaleb, Cherrate, Skhairate and Mansouria stations and in comparison with the control station of Bouznika.

References

1. Adriano, D.C, *Trace Elements in Terrestrial Environnements. Biogeochemistry, Bioavailability and Risks of Metals.* Springer-Verlag, New York (2001).
2. Kabata-Pendias, A., Pendias, H., *Trace Elements in Soils and Plants.* CRS Press, Boca Raton (2001).
3. Nicholson, F.A., Smith, S.R., Alloway, B.J., Carlton-Smith, C., Chambers, B.J., *An inventory of heavy metals inputs to agricultural soils in England and Wales.* *Sci. Total Environ*, 311 (2003) 205–219.
4. CCME, Canadian soil quality guidelines for the protection of environmental and human health: summary tables (2007). In: Canadian Environmental Quality Guidelines, 1999. CCME, Winnipeg.
5. Kabata-Pendias, A.B. Mukherjee., *Trace Elements from Soil to Human.* Springer- Verlag, Berlin (2007).
6. Li, J.L., He, M., Han, W., Gu, Y.F., *Analysis and assessment on heavy metal sources in the coastal soils developed from alluvial deposits using multivariate statistical methods.* *Journal of Hazardous Materials* (2009b). 164, 976–981.
7. Facchinelli, A., Sacchi, E., Mallen, L., *Multivariate statistical and GISbased approach to identify heavy metal sources in soils.* *Environmental Pollution* 114 (2001) 313–324.
8. Sauvé S., McBride M.B., Hendershot W.H., 1997a. *Speciation of lead in contaminated soils.* *Environ. Pollut.* (1998) 149–155.
9. Sauvé, S., Hendershot, W., Allen, H.E., 2000. *Solid-solution partitioning of metals in contaminated soils: dependence on pH, total metal burden, and organic matter.* *Environ. Sci. Technol.* 34, 1125–1131.
10. Matos, A.T., Fontes, M.P.F., da Costa, L.M., Martinez, M.A., *Mobility of heavy metals as related to soil chemicals and mineralogical characteristics of Brazilian soils.* *Environ. Pollut.* (2001). 111, 429–435.

11. Oste, L.A., Temminghoff, E.J.M., Van Riemsdijk, W.H., *Solid-solution partitioning of organic matter in soils as influenced by an increase in pH or Ca concentration. Environ. Sci. Technol.* (2002). 36, 208–214.
12. Hernandez, L., Probst, A., Probst, J.L., Ulrich, E., *Heavy metal distribution in some French forest soils: evidence for atmospheric contamination. Sci. Total Environ.* (2003). 312, 195–219.
13. Micó, C., Recatalá, L., Peris, M., Sánchez, J., *Assessing heavy metal sources in agricultural soils of an European Mediterranean area by multivariate analysis. Chemosphere* (2006). 65, 863–872.
14. Dragovic, S., Mihailovic, N., Gajic, B., *Heavy metals in soils: distribution, relationship with soil characteristics and radionuclides and multivariate assessment of contamination sources. Chemosphere* (2008). 72, 491–495.
15. Amami, B., Serge, D.M., Rhazi, L., Rhazi, M., Bouahim, S., *'Modern pollen–vegetation relationships within a small Mediterranean temporary pool (western Morocco)'*, (2010) *Review of Palaeobotany and Palynology*.
16. El Morhit, M., Fekhaoui, M., El Abidi, A., Yahyaoui, A., *Contamination des muscles de cinq espèces de poissons de l'estuaire du bas Loukkoset qualité de l'alimentation animale au Maroc (côte atlantique marocaine). ScienceLib.*, (2012). 4, 1–21.
17. AFNOR. NF X31-107 ., *Qualité du sol –Détermination de la distribution granulométrique des particules du sol - Méthode à la pipette. Thème: Propriétés physiques des sols* (2003).
18. Walkley, A. and Black, I.A., *An examination of the Degtjareff method for determining organic carbon in soils: Effect of variations in digestion conditions and of inorganic soil constituents* (1934). *Soil Sci.* 63:251-263
19. ISO 11265:1994 *'Qualité du sol'* Détermination de la conductivité électrique spécifique.
20. Mc Lean, E.O., *pH and lime requirements. In: Page, A.L. et al. (Eds.), Methods of Soil Analysis, Part 2, second ed., Agronomy, vol. 9 Soil Society of America, Madison* (1982), WI, pp. 199–244.
21. Méthode de Bernard décrite par Chamley., *Guide Des techniques du laboratoire de Géologie Marine de Luminy*, (1966). 198p.
22. Brallier, S., Harrison, R.B., Henry, C.L., Dongsen, X., *Liming effects on availability of Cd, Cu, Ni and Zn in a soil amended with sewage sludge 16 years previously. Water, Air and soil Pollution*, (1996). 86, 195-206.
23. Baize, D., *Teneurs totales en éléments traces métalliques dans les sols (France). Reference set strategies d'interprétation. INRA Editions, Paris.* (1997). 410 p.
24. Navel, A., *Distribution, spéciation, impact et transfert du cuivre dans un sol sous vigne: rôle de la structuration spatiale et du statut organique Thèse de Doctorat, l'Université de Grenoble* (2011), 253P.
25. Godin, *sources de contaminations et enjeux. Séminaire: «Eléments traces et pollution des sols»*, 4-5 mai 1982 Paris, PP 3-12.
26. El Morhit, M., Fekhaoui, M., Elie, P., Girard, P., Yahyaoui, A., ElAbidi, A., Jbilou, M., *Heavy metals in sediment, water and the European glass eel, Anguilla Anguilla (Osteichthyes :Anguillidae) from Loukkos river estuary (Morocco eastern Atlantic). Cybium*, (2009). 33 (3), 219–228.
27. Lee, C.G., Chon, H.T., Jung, M.C., *Heavy metal contamination in the vicinity of the daduk Au-Ag-Pb-Zn mine in Korea. Applied Geochemistry*, (2001). 16: 1377-1386.
28. Chon, H.T., Ahn, J.S., Jung, M.C., *Seasonal variations and chemical forms of heavy metals in soils and dusts from the satellite cities of Seoul, Korea. Environ. Geochem. Health* (1998). 20, 77–86.
29. Nimick, D.A & Moor, J.M., *Prediction of water-soluble metal concentrations in fluviually deposited tailing sediments, Upper Clarck Fotk Valley, Montana, USA. Applied Geochem*, (1991). 6: 635-646.
30. Jung, M.C., *Environnemental contamination of heavy metals in soils, plans, waters and sediments in the vicinity of metaliferous mine in Korea. Unpubl. PhD thesis, Univ, London*, (1995). pp: 189-317.

(2014) ; <http://www.jmaterenvirosci.com>